

Technical Alert

Sexually Transmitted Infection Testing in Young Patients

TO: Medical Staff and Clients

FROM: Dr. Amy Woron, PhD, MPH

V.P. - Technical Director (Microbiology/Molecular)

Jason Pon MB(ASCP), CHS(ABHI) Dr. Wesley Kim, MD Dr. Ana Ortega-Lopez, MD

Manager, DLS Molecular Lab Medical Director DLS Medical Director QMC Punchbowl,

and QMC West North Hawaii and Molokai

DATE: Revalidated May 2, 2024

SUBJECT: Testing for Chlamydia and gonorrhea is performed at a reference laboratory for patients under

the age of 14 years

Although detection of *Chlamydia trachomatis* (CT) and *Neisseria gonorrhoeae* (GC) in urogenital specimens by nucleic acid amplification test (NAAT) is available at DLS, the FDA-labeling of this product is age-restricted to patients who are at least 14 years old. Neither is it labeled for the evaluation of suspected abuse cases. The test of choice in these circumstances is culture because of specificity, although this approach lacks sensitivity and collection methods are more invasive. The Centers for Disease Control and Prevention is also very cautious in its recommendations for testing young patients (https://www.cdc.gov/mmwr/preview/mmwrhtml/rr6302a1.htm).

DLS has also secured a commitment from a reference laboratory to perform this off-label NAAT testing for our customers. Alternatively, there is the availability of CT/GC culture capability (order codes 6201 and 625, respectively). The send-out order code (5127) and specimen collection devices (Aptima®) are different from the inhouse order code (183) and our Cobas collection products. The Aptima® devices are available from Client Services.

These tests are made available for the expressed purpose of medical management of the patient and are not intended or appropriate for law enforcement or investigative purposes.

Order code: 5127

Test name: CT/GC RNA, TMA, Urogenital

Preferred specimen type: Urine

Preferred container: Sterile Container, minimum volume 2 mL

Special instructions: Patient should not have urinated within one hour prior to collection. Female patients

should not cleanse the labial area prior to providing the specimen. Direct patient to provide a first-catch urine (a maximum of 20-30 mL of the initial urine stream) into

a urine collection cup free of any preservatives.

Other acceptable types: 0.5mL of SurePathTM preservative fluid collected in APTIMA® specimen transfer

Tube (Green label) or APTIMA® Vaginal collection or Multitest collection kit (Orange label), or 2mL urine using the APTIMA® urine specimen collection kit.

Specimen stability (in APTIMA®): Ambient: 30 days, Refrigerated: 30 days, Frozen: 1 year

Specimen stability (not in APTIMA®): Refrigerated: 24 hours

Please refer any questions to Jason Pon, Manager - DLS Molecular Laboratory at 808-441-5469 or DLS Client Services at 808-589-5101.